

For adults with Chronic Idiopathic Constipation (CIC)*

Here's your completed

Doctor Discussion Guide

Share this with your doctor to make the conversation about Chronic Constipation a little bit easier

CIC can be difficult to put into words. Let this guide help you and your doctor talk about the issues specific to your CIC experience. If you're looking to change your current approach, ask your doctor about the different ways to treat CIC.

*Chronic means having symptoms for at least 6 months. Idiopathic means the cause of the constipation is unknown.

IMPORTANT SAFETY INFORMATION

Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

Scroll below for additional Important Safety Information and the full Prescribing Information and Patient Information.



Treatment History

1

What treatments are you currently taking, if any?

2

What treatments have you tried in the past, if any?

3

Are you still taking any of those treatments? If not, how long did you take those treatments?

4

For past treatments, why did you stop taking those medications?



Personal Life

5

Does your CIC affect how you are feeling?

6

How often are your daily activities impacted by your CIC?

7

What symptoms are bothering you?

8

How often do your symptoms bother you?



Healthy Lifestyle

9

Are you following a high fiber diet?

10

Do you make an effort to drink more water and stay hydrated?

Scroll below for additional Important Safety Information and the full Prescribing Information and Patient Information.



Bathroom Visits

11

Do your visits to the toilet leave you feeling like you have not fully finished?

12

Do you often plan your day around trips to the bathroom?

13

Does going to the toilet ever disrupt your daily routine?

14

Do you find yourself looking for the nearest bathroom wherever you are “just in case”?

15

How many days a week does CIC typically disrupt your day?

You're all set to

talk to your doctor!

Hopefully, this guide will make it easier to discuss the symptoms you have been experiencing, and to help determine a treatment plan for you.


Scroll below for additional Important Safety Information and the full Prescribing Information and Patient Information.

Make your move with Motegrity,

the only CIC treatment of its kind


Motegrity helps get your colon moving

Unlike other prescription medications for CIC, Motegrity works differently by stimulating natural movements of the colon muscle, known as peristalsis.




Ready

Motegrity is absorbed by your body.



Set

Motegrity gets to work by stimulating the natural muscle movements of your colon.



Move

This all works to help make a bowel movement.

**Ask your doctor if
Motegrity may be
right for you.**

IMPORTANT SAFETY INFORMATION

Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

Before taking Motegrity, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have or have had depression, suicidal thoughts or actions, or mood problems
- have kidney problems. Your HCP may give you a lower dose of Motegrity
- are pregnant or plan to become pregnant. It is not known if Motegrity will harm your unborn baby
- are breastfeeding or plan to breastfeed. Prucalopride can pass into your breastmilk. Talk with your HCP about the best way to feed your baby if you take Motegrity.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of Motegrity? Motegrity may cause serious side effects, including:

- unusual changes in mood or behavior, thoughts of hurting yourself, trying to hurt yourself, or suicide. Stop taking Motegrity right away and tell your HCP immediately if your depression gets worse, you feel sad, hopeless or begin to have thoughts of suicide, thoughts of hurting yourself or you have tried to hurt yourself or if you develop new depression.

The most common side effects of Motegrity include:

- headache
- stomach area (abdominal) pain or bloating
- nausea
- diarrhea
- dizziness
- vomiting
- gas
- fatigue

These are not all the possible side effects of Motegrity. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

What is Motegrity® (prucalopride)?

Motegrity is a prescription medicine used in adults to treat a type of constipation called chronic idiopathic constipation (CIC). Idiopathic means the cause of the constipation is unknown. It is not known if Motegrity is safe and effective in children.

Please click here for full [Prescribing Information](#) and [Patient Information](#).



©2023 Takeda Pharmaceuticals U.S.A., Inc., 95 Hayden Ave., Lexington, MA 02421. All rights reserved. 1-877-TAKEDA-7 (1-877-825-3327). MOTEGRITY® and the MOTEGRITY Logo® are registered trademarks of Takeda Pharmaceuticals U.S.A., Inc. TAKEDA and the TAKEDA logo are registered trademarks of Takeda Pharmaceutical Company Limited. US-PRU-0375v2.0 01/23

motegrity®
(prucalopride) tablets 1mg, 2mg