

**Maximize savings**

with a 90-day supply

See page 11 for details.

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

I chose the one that  
**m**oves me



*Actor Portrayal  
Individual results may vary*

for adults with **Chronic  
Idiopathic Constipation (CIC)\***

**What to expect with Motegrity**—by increasing rhythmic movements of the colon muscle, it is the **only prescription medicine of its kind** that works differently for adults with CIC.

\*The ability of Motegrity to relieve the infrequency of bowel movements (BMs) was studied in six clinical trials lasting 12 weeks (Studies 1-5) and 24 weeks (Study 6), included 2,484 adults with CIC. About 3 in 4 were women and white, with an average age of 47 (ranging from 17 to 95), and many had been living with symptoms of CIC for years. During 12-week studies, Motegrity was shown to help normalize the number of complete spontaneous bowel movements (CSBMs) per week for adults living with CIC (at least 3 CSBMs per week on average). Chronic means having symptoms for at least 6 months. Idiopathic means the cause of the constipation is unknown.

## IMPORTANT SAFETY INFORMATION

### Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

Please see complete Important Safety Information on page 10 and [Prescribing Information](#).

Remember to check in with your doctor  
and tell them how you're doing.

CHECK IN BY

PHONE NUMBER

\_\_\_/\_\_\_ (\_\_\_) \_\_\_-\_\_\_

# Congratulations on your move to Motegrity

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg



## It works differently by moving your colon muscle to help normalize complete spontaneous bowel movements (CSBMs)

- Unlike other prescription medications for CIC, Motegrity enhances the **natural movements of the colon muscle**, known as peristalsis
- Motegrity is classified as a “prokinetic” because it works by promoting **colonic movements**
- There are many prescription therapies for treating Chronic Idiopathic Constipation (CIC), but only one that works like **Motegrity**

Chronic means having symptoms for at least 6 months. Idiopathic means the cause of constipation is unknown.

## IMPORTANT SAFETY INFORMATION

**Before taking Motegrity, tell your healthcare provider (HCP) about all of your medical conditions, including if you:**

- have or have had depression, suicidal thoughts or actions, or mood problems
- have kidney problems. Your HCP may give you a lower dose of Motegrity
- are pregnant or plan to become pregnant. It is not known if Motegrity will harm your unborn baby
- are breastfeeding or plan to breastfeed. Prucalopride can pass into your breastmilk. Talk with your HCP about the best way to feed your baby if you take Motegrity.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.


**Please see complete Important Safety Information on page 10 and [Prescribing Information](#).**

# How Motegrity helps move the colon muscle

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

Motegrity is the only prescription medicine of its kind for adults with CIC. Check out the diagram below to learn more about how Motegrity is designed to move your colon muscle.

## Motegrity works in three steps to help get you moving



### 1 Ready

Taking Motegrity releases an active ingredient that gets absorbed into your system.

### 2 Set

It targets a specific receptor\* in your colon and enhances the natural movements of the colon muscle.

### 3 Move

Rhythmic muscle movements in your colon are increased. This helps your bowels move.

\*Receptors are molecules that help cells communicate and bring about responses, such as colon muscle movements.

## IMPORTANT SAFETY INFORMATION

**What are the possible side effects of Motegrity?**

**Motegrity may cause serious side effects, including:**

- unusual changes in mood or behavior, thoughts of hurting yourself, trying to hurt yourself, or suicide. Stop taking Motegrity right away and tell your HCP immediately if your depression gets worse, you feel sad, hopeless or begin to have thoughts of suicide, thoughts of hurting yourself or you have tried to hurt yourself or if you develop new depression.

Please see complete Important Safety Information on page 10 and [Prescribing Information](#).

# Motegrity moved patients to share their stories

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## Hear from real patients who spoke to their doctor about how Motegrity could move them

CIC is a chronic bowel condition affecting as many as 17.8 million adults in the United States.\* You are not alone in this. You can be open when discussing your symptoms with your doctor and talk about how your Motegrity treatment is going.

\*Based on US prevalence of 6.9%.



I've worked really closely with my GI doctor to manage my symptoms and figure out a new normal for me.

**- Terri**

Real Motegrity patient



Being my own best advocate helped me find a CIC treatment option that worked for me.

**- Matt**

Real Motegrity patient



See their full videos at [motegrity.com/patient-stories](https://motegrity.com/patient-stories)

## IMPORTANT SAFETY INFORMATION

### The most common side effects of Motegrity include:

- headache
- stomach area (abdominal) pain or bloating
- nausea
- diarrhea
- dizziness
- vomiting
- gas
- fatigue

These are not all the possible side effects of Motegrity. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see complete Important Safety Information on page 10 and [Prescribing Information](#).

# Motegrity moved patients to share their stories

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg



After discussing my diagnosis with my doctor, we decided Motegrity may be an appropriate treatment for me. It was different than other treatments I had tried, and it made a difference for me.

**- Lauren**  
Real Motegrity patient



I feel relief from my CIC and I am able to go to the bathroom more often. I don't feel like I'm chained to a toilet.

**- Kerry**  
Real Motegrity patient



See their full videos at [motegrity.com/patient-stories](https://motegrity.com/patient-stories)

## IMPORTANT SAFETY INFORMATION

### Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

Please see complete Important Safety Information on page 10 and [Prescribing Information](#).

# Here's what to expect with Motegrity

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## A different way to help you achieve CIC relief

The ability of Motegrity to relieve the infrequency of bowel movements was studied in six clinical trials with 2,484 adults living with Chronic Idiopathic Constipation (CIC).\*



While your results may vary, here's how Motegrity was shown to help:

### Motegrity may help you get that “complete” feeling more often

If you've lived with CIC, you may know how rare getting those “complete” moments can be when it comes to your bowel movements (BMs).

During 12-week studies, Motegrity was shown to help **normalize** the number of complete spontaneous BMs per week for adults living with CIC.

Normalize means at least 3 complete spontaneous BMs per week on average.

\*In the studies, about 3 in 4 were women and white, with an average age of 47 (ranging from 17 to 95), and many had been living with symptoms of CIC for years.

## IMPORTANT SAFETY INFORMATION

**Before taking Motegrity, tell your healthcare provider (HCP) about all of your medical conditions, including if you:**

- have or have had depression, suicidal thoughts or actions, or mood problems
- have kidney problems. Your HCP may give you a lower dose of Motegrity
- are pregnant or plan to become pregnant. It is not known if Motegrity will harm your unborn baby
- are breastfeeding or plan to breastfeed. Prucalopride can pass into your breastmilk. Talk with your HCP about the best way to feed your baby if you take Motegrity.

**Please see complete Important Safety Information on page 10 and [Prescribing Information](#).**

# Motegrity helps get you moving

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## CIC relief held steady through 12 weeks<sup>†</sup>

In clinical studies, Motegrity improved the number of complete spontaneous BMs per week as early as Week 1 and lasted 3 months (the length of the studies).

<sup>†</sup>Motegrity was evaluated in studies lasting up to 12 weeks.

## People got that “complete” moment sooner with Motegrity

People who started taking Motegrity had their first complete spontaneous BM 1.4 to 4.7 days into treatment.<sup>‡</sup>

<sup>‡</sup>Compared to patients taking a placebo—a pill with no active ingredient—ranged from 1.4 to 4.7 days compared with 9.1 to 20.6 days across 6 studies.



## Notes:

---

---

---

---

## IMPORTANT SAFETY INFORMATION

**What are the possible side effects of Motegrity?**  
**Motegrity may cause serious side effects, including:**

- unusual changes in mood or behavior, thoughts of hurting yourself, trying to hurt yourself, or suicide. Stop taking Motegrity right away and tell your HCP immediately if your depression gets worse, you feel sad, hopeless or begin to have thoughts of suicide, thoughts of hurting yourself or you have tried to hurt yourself or if you develop new depression.

**Please see complete Important Safety Information on page 10 and [Prescribing Information](#).**

# Motegrity was extensively studied

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## Motegrity was evaluated across 6 studies



The safety profile of Motegrity was studied in six clinical trials involving more than 2,500 adults with CIC, about half of whom received Motegrity.

## Motegrity may cause serious side effects, including:



unusual changes in mood or behavior, thoughts of hurting yourself, trying to hurt yourself, or suicide. Stop taking Motegrity right away and tell your healthcare provider immediately if your depression gets worse, you feel sad, hopeless or begin to have thoughts of suicide, thoughts of hurting yourself or you have tried to hurt yourself or if you develop new depression.

## What are the common side effects for Motegrity?

**The most common side effects of Motegrity include:** headache, stomach area (abdominal) pain or bloating, nausea, diarrhea, dizziness, vomiting, gas, and fatigue.

These are not all the possible side effects of Motegrity. Call your doctor for medical advice about side effects.

For patients who experienced diarrhea or headache, those side effects typically occurred within the first week of treatment and resolved within a few days of continued use.

**Talk to your doctor to learn more about Motegrity.**



Please see complete Important Safety Information on page 10 and [Prescribing Information](#).



# Once-daily dosing

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## Convenient and simple dosing for your busy schedule

There's no need to complicate things with Motegrity. Just one tablet a day. Anytime. With or without food. And of course, always follow your doctor's specific instructions. If you have kidney problems, your doctor may give you a lower dose.



**1 tablet a day**



**Anytime**



**With or without food**



## Staying on track can help you get the most out of treatment

Motegrity was shown to be effective when taken once daily, so it's very important to take it every day as prescribed—not just on days when you are experiencing symptoms.



### Healthy lifestyle tips for CIC

- Maintain a high-fiber diet
- Drink plenty of fluids
- Participate in regular physical activity

## IMPORTANT SAFETY INFORMATION

### Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

**Please see complete Important Safety Information on page 10 and [Prescribing Information](#).**

## IMPORTANT SAFETY INFORMATION

### Do not take Motegrity if you:

- are allergic to Motegrity. Allergic reaction symptoms may include trouble breathing, rash, itching and swelling of your face, lips, tongue or throat.
- have a tear in your stomach or intestinal wall (bowel perforation), a bowel blockage (intestinal obstruction) or serious conditions of the intestinal wall such as Crohn's disease, ulcerative colitis.

### Before taking Motegrity, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have or have had depression, suicidal thoughts or actions, or mood problems
- have kidney problems. Your HCP may give you a lower dose of Motegrity
- are pregnant or plan to become pregnant. It is not known if Motegrity will harm your unborn baby
- are breastfeeding or plan to breastfeed. Prucalopride can pass into your breastmilk. Talk with your HCP about the best way to feed your baby if you take Motegrity.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

### What are the possible side effects of Motegrity?

#### Motegrity may cause serious side effects, including:

- unusual changes in mood or behavior, thoughts of hurting yourself, trying to hurt yourself, or suicide. Stop taking Motegrity right away and tell your HCP immediately if your depression gets worse, you feel sad, hopeless or begin to have thoughts of suicide, thoughts of hurting yourself or you have tried to hurt yourself or if you develop new depression.

#### The most common side effects of Motegrity include:

- headache
- dizziness
- stomach area (abdominal) pain or bloating
- vomiting
- nausea
- gas
- diarrhea
- fatigue

These are not all the possible side effects of Motegrity. Call your doctor for medical advice about side effects.

**You are encouraged to report negative side effects of prescription drugs to the FDA.**

Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

## What Is Motegrity<sup>®</sup> (prucalopride)?

Motegrity is a prescription medicine used in adults to treat a type of constipation called chronic idiopathic constipation (CIC). Idiopathic means the cause of the constipation is unknown. It is not known if Motegrity is safe and effective in children.

Please click here for full [Prescribing Information](#) and [Patient Information](#).

# Maximize savings with a 90-day supply

**motegrity**<sup>®</sup>  
(prucalopride) tablets 1mg, 2mg

## Pay just \$15 for a 3-month supply

With the free Motegrity Savings Card, appropriate, eligible patients pay as little as **\$15 for a 30- or 90-day prescription fill\***

## Two simple ways to sign up for your Motegrity Savings Card



1. Text **SAVE2DAY** to 36395<sup>†</sup> to sign up and receive refill reminders via text.
2. Visit [Motegrity.com/signup](https://www.motegrity.com/signup).

Now you can keep your Motegrity Savings Card on your smartphone. Simply add to your Mobile Wallet and use at the pharmacy.



\*Up to \$325 max benefit per use per 90-day prescription; \$90 max benefit per use per 30-day prescription. See Terms and Conditions below.

†Message & data rates may apply. Average 5 messages per month. Text HELP to 36395 for info, STOP to end. View terms and conditions: [Motegrity.com/signup](https://www.motegrity.com/signup).

**Terms & Conditions:** Eligible Commercially Insured patients may pay as little as \$15 and receive up to \$90 off their co-pay or out of pocket expenses per 30 day supply of Motegrity<sup>®</sup> (prucalopride). Offer is tiered based on quantity dispensed: Tier 1: 1-30 tablets; Patient pays \$15, up to max \$90 benefit for 1 use (\$2700 lifetime); Tier 2: 31-60 tablets; Patient pays \$15, up to max \$180 benefit for 2 use(s) (\$2700 lifetime); Tier 3: 61-90 tablets; Patient pays \$15, up to max \$325 benefit for 3 use(s) (\$3250 lifetime). Offer valid for up to 30 uses through offer expiry. A valid Prescriber ID# is required on the prescription. Offer valid for patients ages 18 and older. Offer not valid for cash paying patients.

**Restrictions:** This offer is valid in the United States, Puerto Rico, and the U.S. Territories. Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, Tricare, or other federal or state health programs (including any state medical assistance program) or where prohibited by the health insurance provider. Cash Discount Cards and other non-insurance plans are not valid as primary under this offer. If the patient is eligible for drug benefits under any such program, the patient cannot use this offer. By using this offer, the patient certifies that he or she will comply with any terms of his or her health insurance contract requiring notification to his or her payor of the existence and/or value of this offer. Offer not valid for patients under 18 years of age. It is illegal to (or offer to) sell, purchase, or trade this offer. This offer is not transferable and is limited to one offer per person and may not be combined with any other coupon, discount, prescription savings card, rebate, free trial, patient assistance, or other offer. Not valid if reproduced. Void where prohibited by law. Program managed by ConnectiveRx on behalf of Takeda Pharmaceuticals U.S.A., Inc. The parties reserve the right to discontinue, rescind, revoke or amend this offer without notice at any time. This is not health insurance.

## IMPORTANT SAFETY INFORMATION

**Before taking Motegrity, tell your healthcare provider (HCP) about all of your medical conditions, including if you:**

- have or have had depression, suicidal thoughts or actions, or mood problems
- have kidney problems. Your HCP may give you a lower dose of Motegrity
- are pregnant or plan to become pregnant. It is not known if Motegrity will harm your unborn baby
- are breastfeeding or plan to breastfeed. Prucalopride can pass into your breastmilk. Talk with your HCP about the best way to feed your baby if you take Motegrity.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Please see complete Important Safety Information on page 10 and [Prescribing Information](#).**



©2023 Takeda Pharmaceuticals U.S.A., Inc., 300 Shire Way, Lexington, MA 02421. 1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. TAKEDA<sup>®</sup> and the TAKEDA Logo<sup>®</sup> are registered trademarks of Takeda Pharmaceutical Company Limited. MOTTEGRITY<sup>®</sup> and the MOTTEGRITY Logo<sup>®</sup> are registered trademarks of Takeda Pharmaceuticals U.S.A., Inc. US-PRU-0365v6.0 11/23